

On the growth of modular symbols

Anton Deitmar

Introduction

Let f be a holomorphic cusp form of weight two for the group $\Gamma = \Gamma_0(N)$. Then $f(z)dz$ is a Γ -invariant holomorphic differential on the upper half plane $\mathbb{H}$ in $\mathbb{C}$. For $\gamma \in \Gamma$ define the modular symbol

$$\langle \gamma, f \rangle \stackrel{\text{def}}{=} -2\pi i \int_{z_0}^{\gamma z_0} f(z) dz,$$

which is independent of the choice of the point $z_0 \in \mathbb{H} \cup \mathbb{Q} \cup \{i\infty\}$. The modular symbol for fixed f is a group homomorphism from Γ to the additive group of complex numbers. For $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ let $\|\gamma\| = \max(|a|, |b|, |c|, |d|)$. In this paper we show that for given f the modular symbol has logarithmic growth, i.e.,

$$|\langle \gamma, f \rangle| \leq A \log \|\gamma\| + B$$

for some $A, B \geq 0$. In [2, 3], D. Goldfeld conjectured that the modular symbol has moderate growth if also f is allowed to vary among the normalized newforms. The result of the present paper reduces Goldfeld's conjecture to a statement on the growth of the modular symbol on a set of generators of the group $\Gamma_0(N)$.

Note that, as f is a cusp form, the modular symbol vanishes on parabolic elements, that is, $\langle p, f \rangle = 0$ for every parabolic element p of Γ .

1 Growth of an additive homomorphism

Let $\mathrm{SL}_2(\mathbb{R})$ denote the group of real 2×2 matrices of determinant one. Let G be the group $\mathrm{PSL}_2(\mathbb{R}) = \mathrm{SL}_2(\mathbb{R}) / \pm 1$.

For $x = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in G$ let $\|x\| = \max(|a|, |b|, |c|, |d|)$. Let M be a subset of G . A function $f : M \rightarrow \mathbb{C}$ is said to be of *logarithmic growth*, if there are constants $A, B \geq 0$ such that

$$|f(x)| \leq A \log \|x\| + B$$

holds for every $x \in M$.

Theorem 1.1 *Let Γ be a lattice in $\mathrm{PSL}_2(\mathbb{R})$ and let $\psi : \Gamma \rightarrow \mathbb{C}$ be a group homomorphism with $\psi(p) = 0$ for every parabolic element of Γ . Then ψ is of logarithmic growth.*

Proof: Let $\mathbb{H}$ denote the upper half plane in $\mathbb{C}$ equipped with the hyperbolic metric $ds^2 = \frac{dx^2 + dy^2}{y^2}$. Choose $z_0 \in \mathbb{H}$ which is not a fixed point of an elliptic element of Γ , and let $\mathcal{F}$ denote the corresponding *Dirichlet fundamental domain*, also called the *Dirichlet polygon* [1], as it is a hyperbolic polygon with finitely many sides. It is defined as

$$\mathcal{F} = \{z \in \mathbb{H} : d(z, z_0) < d(\gamma z, z_0) \ \forall \gamma \in \Gamma \setminus \{1\}\},$$

where $d(z, w)$ denotes the hyperbolic distance of two points in the upper half plane $\mathbb{H}$. Let $S = S^{-1}$ be a finite set of generators of the group Γ . For $\gamma \in \Gamma$ we write $\gamma = s_1 \cdots s_n$ as a shortest word, so $n = l_S(\gamma)$, the *word length* of γ with respect to S . Then,

$$|\psi(\gamma)| = |\psi(s_1 \cdots s_n)| = \left| \sum_{j=1}^n \psi(s_j) \right| \leq C_S l_S(\gamma),$$

where $C_S \geq 0$ is the maximum of the values $|\psi(s)|$ for $s \in S$.

We first consider the case of a uniform lattice Γ , i.e., the quotient $\Gamma \backslash G$ is compact. Then the closure $\bar{\mathcal{F}}$ of $\mathcal{F}$ in $\mathbb{H}$ is compact. According to Theorem IV

23 of [4], for every $z \in \mathbb{H}$ there exists $\lambda \geq 1$, $C \geq 0$ with $l_S(\gamma) \leq \lambda d(\gamma z, z) + C$ for every $\gamma \in \Gamma$. For $z = i$ this implies

$$|\psi(\gamma)| \leq C_S \lambda d(\gamma i, i) + C_S C.$$

By Theorem 7.2.1 of [1] one has for $z, w \in \mathbb{H}$,

$$d(z, w) = \log \left(\frac{|z - \bar{w}| + |z - w|}{|z - \bar{w}| - |z - w|} \right).$$

Let $\gamma = \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$ we get

$$\begin{aligned} d(\gamma i, i) &= \log \left(\frac{\left| \frac{ai+b}{ci+d} + i \right| + \left| \frac{ai+b}{ci+d} - i \right|}{\left| \frac{ai+b}{ci+d} + i \right| - \left| \frac{ai+b}{ci+d} - i \right|} \right) \\ &= \log \left(\frac{|ai + b - c + di| + |ai + b + c - di|}{|ai + b - c + di| - |ai + b + c - di|} \right) \\ &= \log \left(\frac{\sqrt{(b-c)^2 + (a+d)^2} + \sqrt{(b+c)^2 + (a-d)^2}}{\sqrt{(b-c)^2 + (a+d)^2} - \sqrt{(b+c)^2 + (a-d)^2}} \right). \end{aligned}$$

Now $(b-c)^2 + (a+d)^2 - ((b+c)^2 + (a-d)^2) = -4bc + 4ad = 4$, as the determinant of γ is 1. Therefore,

$$\begin{aligned} d(\gamma i, i) &= \log \left(\left(\sqrt{(b-c)^2 + (a+d)^2} + \sqrt{(b+c)^2 + (a-d)^2} \right)^2 \right) - 2 \log 2 \\ &\leq 2 \log \left(\sqrt{8 \|\gamma\|^2} + \sqrt{8 \|\gamma\|^2} \right) - 2 \log 2 \\ &= 2 \log \|\gamma\| + 3 \log 2. \end{aligned}$$

This gives the Theorem in the case of Γ being a uniform lattice.

Next assume that Γ is not uniform. Then the fundamental domain $\mathcal{F}$ has cusps. We assume that z_0 is chosen in a way that no two cusps of $\mathcal{F}$ are equivalent under Γ . So the cusps of $\mathcal{F}$ are a set of representatives of the set of cusps of Γ in the boundary $\partial\mathbb{H}$ of $\mathbb{H}$ module Γ -equivalence. For each cusp $c \in \partial\mathbb{H}$ fix some $\sigma_c \in G$ with $c = \sigma_c \infty$. We do so in a Γ -compatible way, i.e., for $\gamma \in \Gamma$ we suppose that $\sigma_{\gamma c} \sigma_c^{-1}$ lies in Γ .

Let $T > 1$ and set

$$\mathcal{F}_T = \{z \in \mathcal{F} : \text{Im}(\sigma_c^{-1}z) \leq T \text{ for every cusp } c\}.$$

We choose T so large that $\mathcal{F}_T$ equals $\mathcal{F}$ minus cusp sections. Let $\mathbb{H}_T$ be the union of all sets $\overline{\gamma\mathcal{F}_T}$ where γ ranges over Γ . Then $\mathbb{H}_T$ equals $\mathbb{H}$ minus a countable number of open horoballs. Thus $\mathbb{H}_T$ is a Riemannian manifold with boundary. Let d_T denote the distance function on $\mathbb{H}_T$. Note that if $z, w \in \mathbb{H}_T$, and the geodesic in $\mathbb{H}$ joining them lies completely in $\mathbb{H}_T$, then $d_T(z, w) = d(z, w)$.

For a cusp c let

$$H_{c,T} = \sigma_c(\{\text{Im}(z) > T\})$$

be the T -horoball attached to c . Increasing T if necessary, we can make sure, that the geodesic $\overline{z_0, \gamma z_0}$ in $\mathbb{H}$ is disjoint to $\mathcal{F} \cap H_{c,T}$ for every cusp c of $\mathcal{F}$ and every $\gamma \in \Gamma$. Note that for every cusp c of Γ , which is not a cusp of $\mathcal{F}$, the intersection $\mathcal{F} \cap H_{c,T}$ is empty.

Let $\gamma \in \Gamma$ and suppose that the geodesic $\overline{z_0, \gamma z_0}$ in $\mathbb{H}$ does not completely lie in $\mathbb{H}_T$. Then this geodesic meets some horoball $H_{c,T}$. After applying σ_c^{-1} , one can assume $c = \infty$. Then there is a generator p_c of the stabilizer group Γ_c of the cusp c , such that the distance $d(z_0, p_c \gamma z_0)$ is then strictly less than $d(z_0, \gamma z_0)$. From this it follows that if $d(z_0, \gamma z_0) \leq d(z_0, p \gamma z_0)$ for every parabolic $p \in \Gamma$, then the geodesic $\overline{z_0, \gamma z_0}$ lies completely in the set $\mathbb{H}_T$.

By Theorem IV 23 in [4], there are $\lambda \geq 1$ and $C \geq 0$ with $l_S(\gamma) \leq \lambda d_T(z_0, \gamma z_0) + C$ for every $\gamma \in \Gamma$.

Lemma 1.2 *Let $\gamma \in \Gamma \setminus \{1\}$ be given. There are parabolic elements $p_1, \dots, p_n$ of Γ such that with $\gamma_s = p_n \cdots p_1 \gamma$ one has $d(z_0, \gamma_s z_0) \leq d(z_0, \gamma z_0)$ and the geodesic $\overline{z_0, \gamma_s z_0}$ lies in $\mathbb{H}_T$.*

Proof: Assume first that for every parabolic element $p \in \Gamma$ the distance $d(z_0, \gamma z_0)$ is less than or equal to $d(z_0, p \gamma z_0)$. Then the geodesic $\overline{z_0, \gamma z_0}$ lies in $\mathbb{H}_T$. We set $\gamma_s = \gamma$ and we are done.

Now if there exists a parabolic $p_1 \in \Gamma$ such that $d(z_0, p_1 \gamma z_0) < d(z_0, \gamma z_0)$ then replace γ with $p_1 \gamma$. After that, either the condition above is satisfied or we find a parabolic p_2 such that $d(z_0, p_2 p_1 \gamma z_0) < d(z_0, p_1 \gamma z_0)$. Iteration yields

a sequence $p_1, p_2, \dots \in \Gamma$. This process terminates, as for a given radius r there are only finitely many Γ -conjugates of z_0 in distance $\leq r$. $\square$

To finish the proof of the theorem let $\gamma \in \Gamma$ and consider γ_s as in the lemma. Then $|\psi(\gamma)| = |\psi(\gamma_s)| \leq C_S l_S(\gamma_s) \leq C_S \lambda d_T(z_0, \gamma_s z_0) + C_S C$ and

$$\begin{aligned} d_T(z_0, \gamma_s z_0) &= d(z_0, \gamma_s z_0) \\ &\leq d(z_0, \gamma z_0) \\ &\leq d(z_0, i) + d(i, \gamma i) + d(\gamma i, \gamma z_0) \\ &\leq 2 \log \|\gamma\| + 3 \log 2 + 2d(z_0, i). \end{aligned}$$

$\square$

The value of these results with respect to the Goldfeld conjecture hinges on the control over the constants C, C_S, λ as the group Γ shrinks. The constant C_S depends on the group Γ and on the homomorphism ψ , i.e., if ψ is a modular symbol, on the cusp form f . The constants C and λ , however, do not depend on ψ , therefore they are easier to control. The following explicit estimate might be useful.

Let R be the diameter of $\mathcal{F}_T$ and let B be the closed ball in $\mathbb{H}_T$ around z_0 of radius R . Let $S = \{s \in \Gamma : sB \cap B \neq \emptyset\}$. Then $S = S^{-1}$ is a finite set of generators of Γ . Let

$$r = \inf\{d(B, \gamma B) : \gamma \in \Gamma \setminus S\}.$$

Lemma 1.3 *The number r is > 0 and*

$$|\psi(\gamma)| \leq C_S \left(\frac{2}{r} \log \|\gamma\| + \frac{3 \log 2}{r} + 1 \right).$$

Proof: Let $r_T = \inf\{d_T(B, \gamma B) : \gamma \in \Gamma \setminus S\}$. The proof of Theorem IV 23 of [4] together with the proof of our theorem yields

$$|\psi(\gamma)| \leq \frac{C_S}{r_T} (2 \log \|\gamma\| + 3 \log 2 + 2d(z_0, i)) + C_S$$

Now $r \geq r_T$ and the argument for $r_T > 0$ also implies $r > 0$. Further, varying z_0 the distance $d(z_0, i)$ can be chosen arbitrarily small. $\square$

References

- [1] **Beardon, A.F.:** *The geometry of discrete groups*. Graduate Texts in Mathematics, 91. Springer-Verlag, New York, 1983.
- [2] **Goldfeld, D.:** *Modular elliptic curves and Diophantine problems*. Number theory (Banff, AB, 1988), 157–175, de Gruyter, Berlin, 1990.
- [3] **Goldfeld, D.:** *Modular forms, elliptic curves and the ABC-conjecture*. A panorama of number theory or the view from Baker's garden (Zürich, 1999), 128-147, Cambridge Univ. Press, Cambridge, 2002.
- [4] **de la Harpe, P.:** *Topics in geometric group theory*. Chicago Lectures in Mathematics. University of Chicago Press, Chicago, IL, 2000.
- [5] **Shimura, G.:** *On the factors of the jacobian variety of a modular function field*. J. Math. Soc. Japan 25, 523-544 (1973).

Mathematisches Institut
Auf der Morgenstelle 10
72076 Tübingen
Germany
`deitmar@uni-tuebingen.de`

2.2.2008